### For CPL Lab **Use Only**

# **Cadham Provincial Laboratory**

# **General Requisition**



ONLY ONE SPECIMEN TYPE PER REQUISITION

All areas of the requisition must be completed (please print clearly) See back for requisition/specimen instructions

Cadham Provincial Laboratory

Tel: (204) 945-6123

PATIENT INFORMATION	

P.O. Box 8450 Fax: (204) 786-4770 Winnipeg, MB R3C 3Y1 E-mail: cadham@gov.mb.ca Website: www.gov.mb.ca/health/publichealth/cpl		
RELEVANT CLINICAL INFORMATION	PATIENT INFORMATION	
Outbreak Code:	PHIN: MB Health Reg. #	
Reason for Test:    High risk pre-immunitation   Occupational   Other:   Screening   Other:   Screening   Other:   Other:   Occupational   Other:   Other:	Alternate ID: RCMP# Other Provinces/Territories  Military # Other:	
Relevant History:	Uninsured:	
Autopsy Diabetes Food Borne Illness Cancer/Chemotherapy Dialysis Transplant	Date of Birth: YYYYI MM/DD Sex: * Chart/Clinic/Lab#	
Signs and Symptoms:   ☐ Bronchiolitis ☐ Fever ☐ Lymphadenopathy   ☐ Conjunctivitis ☐ Gastrointestinal ☐ Pneumonia	Patient Legal Last Name * First Name *	
☐ Diarrhea ☐ Influenza-Like Illness ☐ Rash☐ Encephalitis ☐ Jaundice	Street or Other (e.g., General Delivery) * Phone #	
Other: Travel/Treatment History:	City/Municipality/First Nations Reserve Postal Code	
	RETURN REPORT TO:	
SPECIMEN INFORMATION Specimen Type: Specimen Source:	Ordering Practitioner Last Name First name To be completed by HCP	
	Facility Ordering TEST	
COPY REPORT TO:  Date/Time:	Facility Address City/Town	
Other Practitioner Last Name of CA CO Coordinator	Postal Code Phone # Secure Fax #	
Facility 490 Hargrave St. Secure Fax# 204-940-2690	After Hours Contact # for Critical Results:	
SEROLOGY	PARASITOLOGY	
Serology Test Panels (see #1 over)  STBBI Panel Post Exposure: Source Panel (1.3) Prenatal Panel Prenatal HIV OPT OUT (2)	☐ Ova & Parasites ☐ Skin Scrapings ☐ Pinworm Examination ☐ Blood Smears ☐ Identification ☐ Other:	
Post Exposure: Exposed Panel (1)	Bacteriology	
HIV (4) HIV 1/2 Ag/Ab Combo Syphilis Screen  Hepatitis	Culture & Sensitivity (C&S)	
HBsAg  HAV IgM (acute HAV infection)  HBcAb (Total)  HBsAg  HBsAb (Immunity)  HCV Ab	Other: Spore/Sterilizer Testing	
	Gonorrhea	
Nucleic Acid (Plasma Only) (5)	Gonorrhea Culture	
☐ WNV PCR ☐ HCV Genotyping	Chlamydia & Gonorrhea Screen (NAAT)	
☐ WNV PCR ☐ HCV Genotyping ☐ HBV PCR/QUANT ☐ HCV PCR/QUANT	Chlamydia & Gonorrhea Screen (NAAT)  Urine (APTIMA Urine Tube/Yellow) Urethra (APTIMA Unisex Swab)	
☐ WNV PCR       ☐ HCV Genotyping         ☐ HBV PCR/QUANT       ☐ HCV PCR/QUANT         Miscellaneous Serology         Acute Immune Status       Acute Immune Status	Chlamydia & Gonorrhea Screen (NAAT)  Urine (APTIMA Urine Tube/Yellow) Cervix (APTIMA Unisex Swab)  Cervix (APTIMA Unisex Swab)  Referral Isolate:	
☐ WNV PCR       ☐ HCV Genotyping         ☐ HBV PCR/QUANT       ☐ HCV PCR/QUANT             Miscellaneous Serology       Acute Immune Status         Measles       ☐ IgM       ☐ IgG         Mumps       ☐ IgM       ☐ IgG         EBV       ☐ IgM       ☐ IgG	Chlamydia & Gonorrhea Screen (NAAT)  Urine (APTIMA Urine Tube/Yellow) Cervix (APTIMA Unisex Swab) Other:	
☐ WNV PCR         ☐ HCV Genotyping           ☐ HBV PCR/QUANT         ☐ HCV PCR/QUANT    Miscellaneous Serology  Acute Immune Status  Acute Immune Statu  Measles ☐ IgM ☐ IgG CMV ☐ IgM ☐ IgG  Mumps ☐ IgM ☐ IgG EBV ☐ IgM ☐ IgG  Rubella ☐ IgM ☐ IgG HSV ☐ IgM ☐ IgG	Chlamydia & Gonorrhea Screen (NAAT)  Urine (APTIMA Urine Tube/Yellow) Cervix (APTIMA Unisex Swab)  Referral Isolate: Identification Isolate Information:  Susceptibility Testing Isolate Information:	
☐ WNV PCR         ☐ HCV Genotyping           ☐ HBV PCR/QUANT         ☐ HCV PCR/QUANT           Miscellaneous Serology	Chlamydia & Gonorrhea Screen (NAAT)  Urine (APTIMA Urine Tube/Yellow)  Cervix (APTIMA Unisex Swab)  Referral Isolate:  Identification Isolate Information:  VIRUS DETECTION (must specify virus requested)	
☐ WNV PCR         ☐ HCV Genotyping           ☐ HBV PCR/QUANT         ☐ HCV PCR/QUANT           Miscellaneous Serology	Chlamydia & Gonorrhea Screen (NAAT)  Urine (APTIMA Urine Tube/Yellow) Cervix (APTIMA Unisex Swab)  Referral Isolate: Identification Isolate Information:  VIRUS DETECTION (must specify virus requested)  Viral Detection	
☐ WNV PCR       ☐ HCV Genotyping         ☐ HBV PCR/QUANT       ☐ HCV PCR/QUANT         Miscellaneous Serology         Acute Immune Status       Acute Immune Status         Measles ☐ IgM ☐ IgG       CMV ☐ IgM ☐ IgG         Mumps ☐ IgM ☐ IgG       EBV ☐ IgM ☐ IgG         Rubella ☐ IgM ☐ IgG       HSV ☐ IgM ☐ IgG         Varicella ☐ IgM ☐ IgG       Parvo B19 ☐ IgM ☐ IgG         WNV ☐ IgM ☐ IgG       WNV ☐ IgM         ☐ Lyme Ab ☐ H. pylori Ab ☐ Mycoplasma pneumoniae IgM	Chlamydia & Gonorrhea Screen (NAAT)  Urine (APTIMA Urine Tube/Yellow)  Cervix (APTIMA Unisex Swab)  Referral Isolate:  Identification Isolate Information:  VIRUS DETECTION (must specify virus requested)	
☐ WNV PCR         ☐ HCV Genotyping           ☐ HBV PCR/QUANT         ☐ HCV PCR/QUANT           Miscellaneous Serology	Chlamydia & Gonorrhea Screen (NAAT)  Urine (APTIMA Urine Tube/Yellow) Cervix (APTIMA Unisex Swab)  Referral Isolate: Identification Isolate Information:  VIRUS DETECTION (must specify virus requested)  Viral Detection	

#### REQUISITION DEMOGRAPHIC INFORMATION

Mandatory Fields: The specimen will not be tested until all mandatory fields (PHIN, Patient Legal Name, Date of Birth, Sex, Practitioner Name and Address, Source/Type) are provided.

Alternative ID: A unique health ID issued by other authorities such as: RCMP, Military, FNIH/ISC, Other Canadian Provinces, Great-West Life, etc. If no PHIN this is a mandatory field.

Sex: M = Male; F = Female; U = Unknown; A = Ambiguous (Transgender)

#### REPORTS

Secure Fax Number: The fax machine must be in a secure location accessible ONLY to persons requiring reports.

**Report Address:** The address where the report(s) will be sent. Complete information including facility name is required to ensure delivery. All reports will be sent by fax unless otherwise indicated.

Copy Report To: This area can only be filled out or authorized by the ordering practitioner and is intended for another practitioner providing care.

#### REQUISITION TEST ORDERING INFORMATION

Outbreak Code: For Infection Control and Public Health Purposes call Outbreak Coordinator (Microbiology Scientist) at 204-945-7473 for code.

Specimen Type: The nature of the specimen (e.g. aspirate, blood, tissue/biopsy, stool, swab, urine, sputum, serum, plasma, CSF, etc.)

Specimen Source: The anatomical location or site (e.g., throat, right leg wound, etc.) from where the specimen was taken.

- 1) Test Panels (1): Prenatal (HBsAg, Rubella IgG, Syphilis, HIV 1/2 Ag/Ab Combo); Serology STBBI (HBsAg, HCV Ab, Syphilis, HIV 1/2 Ag/Ab Combo); Post Exposure Exposed (HBsAg, HCV Ab, HIV 1/2 Ag/Ab Combo, HBsAb); Post Exposure Source (HBsAg, HCV Ab, HCV Ag, HIV 1/2 Ag/Ab Combo).
- 2) HIV Opt Out Box (2): When this box is checked off, HIV antibody testing will not be conducted as part of the panel.
- 3) Post Exposure Panels (3): If T55 protocol is required, list T55 in the "Other" space under Reason for Test on the front of this form. The "Other" space can also be used if this testing is required due to a bite.
- 4) HIV (Retrovirus) (4): For HIV Viral Load and Genotyping, use HIV/Retrovirus Nucleic Acid Testing Requisition (MG5126).
- 5) Nucleic Acid (5): (Viral load) Send 10 mL EDTA (lavender top) whole blood (must be received within 6 hours at CPL) or EDTA plasma (stored at 2-8°C and received within 3 days at CPL). Please record on the front of this requisition the date and time of collection.

## SPECIMEN COLLECTION INFORMATION

Specimen Labelling: Label specimen with patient's full name and PHIN or alternate ID.

Serology Specimen Volume Requirement: 9 mL serum separator tube (gold top, full draw), preferably spun down.

**Gonorrhea Culture:** Culture is the method of choice for detection of Gonorrhea in Eye, Throat, and Rectal specimens. Use a swab in charcoal transport medium for collection. Culture allows antimicrobial susceptibility testing and supports Gonorrhea surveillance activities.

Chlamydia & Gonorrhea Screen (NAAT): The Aptima Combo 2 Nucleic Acid Ampllification Test (NAAT) is used for routine screening and detection of Chlamydia and Gonorrhea.

**Endocervical Swab Specimens:** The cervical swab is the specimen of choice for women. Use the Aptima Unisex Swab Collection Kit. **Urine:** Urine is the specimen of choice for males. It is also recommended for women without a cervix (e.g. due to hysterectomy) as well as women refusing a complete genital exam. Use the Aptima Urine Collection Kit. First void urine recommended – If this is not practicable, the patient should not have urinated for at least one hour prior to sample collection.

Extragenital Specimens: For screening of Eye, Throat and Rectal specimens, use the Aptima Unisex Swab Collection Kit.

NAAT has replaced DFA and culture as the method of choice for detection of Chlamydia. The Aptima Combo 2 Assay is now Health Canada licensed for use with Rectal and Throat specimens, but culture remains the method of choice for detection of Gonorrhea in extragenital specimens.

The following specimens are unsuitable for NAAT and will not be processed:

- 1) Urine or swab specimens not submitted in Aptima collection tubes;
- 2) Aptima tubes submitted with the foil cap missing or pierced;
- 3) Aptima urine tubes submitted with too much or too little liquid, the liquid level must be between the two arrows on the tube;
- 4) Aptima swab tubes submitted with no swab, 2 swabs, the cleaning swab (white shaft) or a swab not provided in the Aptima collection kit.

#### Virus Detection Specimens:

Once a specimen has been appropriately collected and the swab placed in the Universal Viral Transport (UVT) tube, it must be stored and transported to CPL at 4°C (range 2-8°C).